

Empathic Responding in Preschool-Aged Children With Familial Risk for Autism

Nicole M. McDonald , Haley G. Murphy, and Daniel S. Messinger

Individuals with autism spectrum disorder (ASD) show deficits in social and emotional reciprocity, which often include empathic responding. The younger siblings of children with ASD (*high-risk siblings*) are at elevated risk for ASD and for subclinical deficits in social-emotional functioning. Higher levels of empathy in high-risk siblings during the second and third years of life predict fewer ASD symptoms and likelihood of diagnosis. We conducted a multi-method investigation of empathic responding to an examiner's accident in 30 low-risk and 48 high-risk siblings with ($n = 12$) and without ASD outcomes ($n = 36$) at 4–6 years of age. Empathic responding was measured through behavioral observation and parent report. Prosocial behavior did not differ by ASD outcome. Children with ASD exhibited lower levels of personal distress than high-risk and low-risk siblings without ASD. Per parent report, high-risk siblings without ASD demonstrated higher levels of empathic responding than low-risk children, while the ASD group did not differ from children without ASD on this measure. Higher levels of observed empathic concern, but not prosocial behavior, were associated with lower Social Affect scores on the Autism Diagnostic Observation Schedule in high-risk children. Results suggest that ASD diagnosis and symptoms are associated with reduced emotional responsiveness to an adult's distress, but not associated with deficits in prosocial behavior at preschool age. Results do not support the idea that empathic responding is negatively impacted in a broader autism phenotype. Findings extend previous research by suggesting that empathy may be a protective factor in the social-emotional development of children with familial risk for ASD. *Autism Res* 2017, 0: 000–000. © 2017 International Society for Autism Research, Wiley Periodicals, Inc.

Keywords: empathic responding; high-risk siblings; autism spectrum disorder; preschool age; familial risk

Empathy, the ability to recognize and experience the emotions of others, is a key component of social-emotional functioning. Empathy is associated with positive social outcomes, including prosocial behavior [Eisenberg, Eggum, & Giunta, 2010] and social competence [Zhou et al., 2002] at preschool age. Empathic responding is often impaired in individuals with autism spectrum disorder [ASD; Sigman, Kasari, Kwon, & Yirmiya, 1992]. High-risk siblings (i.e., younger siblings of children with ASD) with ASD outcomes show deficits in empathic responding during the first years of life [Hutman et al., 2010; McDonald & Messinger, 2012]. To extend this area of research, we conducted a multimodal investigation of associations between empathic responding and ASD diagnosis and symptoms in high-risk and low-risk siblings at preschool age.

Empathy is often studied via behavioral responses to distress. Infants demonstrate precursors to empathy through crying in response to other infants' distress

[Sagi & Hoffman, 1976]. Personal distress, a self-oriented aversive emotional reaction, is characterized by discomfort after experiencing the emotions of another [Batson, Fultz, & Schoenrade, 1987; Eisenberg et al., 2010]. Personal distress can act as both an early developmental precursor to empathic responses [Hoffman, 2008], as well as an independent response to emotional situations [Batson et al., 1987; Eisenberg & Eggum, 2009]. By 2 years old, children typically transition from self-oriented responses to the capacity for other-oriented concern [Zahn-Waxler, Radke-Yarrow, Wagner, & Chapman, 1992]. While children are capable of limited prosocial responses (e.g., instrumental helping) by 18 months of age [Zahn-Waxler et al., 1992], 30-month-old children can understand others' emotions and engage in helping behaviors based on implicit emotional cues [Svetlova, Nichols, & Brownell, 2010]. By 3 to 5 years, children typically evidence theory of mind and perspective-taking skills [Wellman, Cross, &

From the Department of Psychiatry and Biobehavioral Sciences, Semel Institute for Neuroscience and Human Behavior, University of California, Los Angeles, Los Angeles, CA 90095, USA (N.M.M.); Department of Psychology, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061 (H.G.M.); Departments of Psychology, Pediatrics, Electrical & Computer Engineering, and Music Engineering, University of Miami, Coral Gables, Florida 33146 (D.S.M.)

Received August 31, 2016; accepted for publication May 15, 2017

Address for correspondence and reprints: Nicole M. McDonald, 760 Westwood Plaza, Room A7-424, Los Angeles, CA 90095. E-mail: nmcdonald@mednet.ucla.edu

Published online 00 Month 2017 in Wiley Online Library (wileyonlinelibrary.com)

DOI: 10.1002/aur.1819

© 2017 International Society for Autism Research, Wiley Periodicals, Inc.

Watson, 2001] and regularly display prosocial behaviors [Eisenberg, Fabes, & Spinrad, 2006].

High-risk siblings with ASD show deficits in empathic responding to familiar and unfamiliar adults in distress from 12 to 36 months of age [Hutman et al., 2010; McDonald & Messinger, 2012]. Likewise, higher levels of empathic responding at 24 months are associated with fewer ASD symptoms at 30 months, suggesting a possible protective role of early empathy abilities in young children with familial risk for ASD [McDonald & Messinger, 2012]. It is less clear, however, how deficits in empathic responding present at preschool age—a critical time for social-emotional development [Denham et al., 2003]. This period is characterized by increased exposure to peers, necessitating increasingly sophisticated social-emotional behaviors, such as emotion identification and communication [Denham et al., 2003] and cooperative play [Coplan & Arbeau, 2009]. Empathy is an important factor in this developing social-emotional competence [Eisenberg et al., 1996, 2006].

Earlier work comparing preschool and school-aged children with ASD to children with typical development and developmental disabilities indicated that ASD was associated with deficits in empathic responding to adults in distress, which were not fully accounted for by differences in intellectual ability [Bacon, Fein, Morris, Waterhouse, & Allen, 1998; Dawson et al., 2004; Sigman et al., 1992]. However, more recent findings suggest that empathy deficits in ASD may not be as pervasive as these earlier studies suggested. Rather, empathy deficits in ASD may be particular to the type of empathy (i.e., cognitive vs. emotional), type of responding (e.g., attention to examiner, prosocial behavior), and mode of measurement [e.g., behavioral observation vs. parent-report; Scheeren, Koot, Mundy, Mous, & Begeer, 2013; Deschamps, Been, & Matthys, 2014; Tsang, Gillespie-Lynch, & Hutman, 2016], highlighting the need for further investigation of individual components of empathy using multiple measures of empathic responding.

There is also a lack information on the degree to which empathy deficits are associated with the broader autism phenotype (BAP), which refers to subclinical ASD characteristics in biological relatives of individuals on the spectrum [e.g., Piven, 2001]. Parents of children with ASD demonstrate lower cognitive empathy abilities than adults with low genetic risk for autism, suggesting a possible role of mild empathy deficits in the BAP [Grove, Baillie, Allison, Baron-Cohen, & Hoekstra, 2014]. A study of high-risk siblings indicated mild deficits in parent-reported cognitive empathy and theory of mind abilities in high-risk siblings without ASD, while deficits in parent-reported emotional empathy and attention to an examiner in distress were not apparent [Tsang et al. 2016].

In the current study, we investigated whether 4- to 6-year-old children with and without familial risk for ASD differed in observed and parent-reported empathic responding based on ASD outcome and level of symptomatology. Children were divided into low risk/no ASD, high risk/no ASD, and ASD groups, and covariates including age, gender, and intellectual ability were evaluated. Individual aspects of empathic responding (empathic concern, prosocial behavior, personal distress) were utilized to uniquely assess whether children with ASD show general or more distinct areas of deficit in their response to an adult in distress. We hypothesized that: (1) children with ASD would show lower levels of emotional responsiveness (empathic concern and personal distress) to an adult in distress than children without ASD, (2) parents of children with ASD would report lower levels of empathic responding than parents of children without ASD, and (3) children with familial risk who had higher levels of empathic responding would have fewer ASD symptoms. We did not expect children without ASD to differ based on familial risk status, nor did we expect group differences in prosocial behavior.

Methods

Participants

Participants were part of a longitudinal study examining the development of infants at elevated risk for ASD due to having an older sibling with ASD at the University of Miami. The sample consisted of 78 children with data from an empathic elicitation task and/or parent report of empathy responding, and ASD outcome data at 4–6 years. There were no differences in demographic variables between ASD outcome groups (see Table I). Children with ASD had lower IQs and higher Calibrated Severity Scores [CSS; Hus, Gotham, & Lord, 2014] on the Autism Diagnostic Observation Schedule [ADOS; Lord et al., 2000] than high-risk and low-risk siblings without ASD.

Procedure

Children visited the laboratory with a caregiver at 4–6 years old. They participated in a battery of behavioral tasks, followed by a diagnostic assessment, which included the Wechsler Preschool and Primary Scale of Intelligence, Third Edition [Wechsler, 2002] and ADOS [Lord et al., 2000].

Measures

Empathic responding. Children's empathic responses were measured with an observational task and a parent questionnaire.

Behavior observation. Following parent-child interaction tasks, the parent was escorted from the room by a

Table I. Participant Information by ASD Outcome Group

Demographic variable	Total <i>n</i> (%)	Low risk/no ASD <i>n</i> (%)	High risk/no ASD <i>n</i> (%)	ASD <i>n</i> (%)	<i>p</i>
ASD outcome	78 (100%)	30 (39%)	36 (46%)	12 (15%)	
Gender					.27
Male	47 (60%)	15 (50%)	23 (64%)	9 (75%)	
Female	31 (40%)	15 (50%)	13 (36%)	3 (25%)	
Race/ethnicity					.79
White/Caucasian	30 (38%)	12 (40%)	12 (33%)	6 (50%)	
Black/African-American	2 (3%)	1 (3%)	1 (3%)	0 (0%)	
Hispanic/Latino	32 (41%)	11 (37%)	17 (47%)	4 (33%)	
Asian/Asian-American	2 (3%)	1 (3%)	0 (0%)	1 (8%)	
More than one race/Other	12 (15%)	5 (17%)	6 (17%)	1 (8%)	
Maternal education					.38
High school	3 (4%)	0 (0%)	3 (8%)	0 (0%)	
Some college	3 (4%)	1 (3%)	2 (6%)	0 (0%)	
2-year college	11 (14%)	5 (17%)	5 (14%)	1 (8%)	
4-year college	25 (32%)	11 (38%)	12 (33%)	2 (17%)	
Advanced/professional degree	35 (45%)	12 (41%)	14 (39%)	9 (75%)	
Measure	<i>M</i> (<i>SD</i>)				
Age at visit (months)	61.87 (9.86)	63.00 (11.82)	62.23 (8.58)	57.96 (7.57)	.32
WPPSI-III Full Scale IQ	102.20 (15.07)	107.59 (11.58)	102.06 (13.13)	89.58 (20.46)	<.01
ADOS-CSS	2.47 (2.20)	1.69 (1.14)	1.83 (1.46)	6.25 (2.18)	<.001
SA-CSS	2.87 (2.51)	2.11 (1.75)	2.25 (1.95)	6.50 (2.51)	<.001
RRB-CSS	3.09 (2.96)	1.29 (1.05)	2.92 (2.58)	7.83 (1.70)	<.001

Notes. Maternal education was unavailable for one child in the low risk/no ASD group. ADOS-CSS was missing for one child in the low risk/no ASD group. WPPSI-III Full Scale IQ scores were missing for one child in the low risk/no ASD group and two children in the high risk/no ASD group. For ADOS severity scores: Autism Spectrum cutoff = 4, Autism cutoff = 6 (Gotham et al., 2009). SA-CSS and RRB-CSS calculated based on criteria presented in Hus et al. [2014].

ASD, autism spectrum disorder; WPPSI-III, Wechsler Preschool and Primary Scale of Intelligence, Third Edition; ADOS, Autism Diagnostic Observation Schedule; CSS, Calibrated Severity Score; SA, Social Affect; RRB, Restricted and Repetitive Behaviors.

research associate and told: “One of the examiners will pretend to hurt herself. We want to see how [child] will respond without anyone else in the room.” A second examiner briefly chatted with the child, then picked up a bin of toys and pretended to hurt herself by dropping it on her foot and saying, “Ow, I hurt my foot. Ow, that really hurt.” This portion of the task was focused on eliciting children’s emotional response to distress. After about 10 sec, if the child had not started picking up the toys, the examiner said “Oh, what a mess!” After several seconds, if the child did not start picking up, the examiner began slowly picking up the toys. This portion of the task was focused on eliciting prosocial behavior. The task ended when the toys were all picked up.

Based on previous work [Young, Fox, & Zahn-Waxler, 1999; McDonald & Messinger, 2012], a coding system was developed to measure children’s responses along three dimensions: empathic concern, prosocial behavior, and personal distress. See Table II for a detailed description of the coding system. A research associate blind to ASD risk group and outcome coded each episode. Approximately 20% of the videos were double coded to assess inter-rater reliability. Intraclass correlations indicated high reliability for each dimension (>.85; see Table II).

Parent report. Empathic responding was also assessed with the Empathy and Prosocial Response subscale of

the Conscience Questionnaire [see Kochanska, DeVet, Goldman, Murray, & Putnam, 1994]. This subscale consists of 13 items measuring children’s empathic responding rated by parents on a 7-point Likert scale (e.g., “Will try to comfort or reassure another in distress,” “Is not likely to become upset if a playmate cries,” “Can tell at just a glance how others are feeling”). Internal consistency within the sample was good (Cronbach’s alpha = .80).

ASD outcome. Following the 4–6-year visit, children fell into three outcome groups: low risk/no ASD, high risk/no ASD, and ASD. A licensed psychologist who was blind to risk group made diagnoses according to DSM-IV-TR criteria [American Psychiatric Association, 2000], and informed by results from the Autism Diagnostic Inventory-Revised [administered at 36 months; Lord, Rutter, & Le Couteur, 1994] and ADOS [administered at 4–6 years; Lord et al., 2000]. Within the high-risk group, 12 of 48 (25%) children were diagnosed with ASD. Calibrated Severity Scores for the Social Affect (SA-CSS) and Restricted and Repetitive Behaviors (RRB-CSS) ADOS subdomains were calculated as a measure of ASD symptoms [Hus et al., 2014]. SA-CSS and RRB-CSS ranged from 1 to 10, indicating scores comprising the

Table II. Description of Empathy-Related Dimensions

Measure	Description	Ratings	ICC
Empathic Concern	Facial, gestural, and vocal signs of sympathy and concern	1—no signs of concern; may ignore examiner, show inappropriate affect, or otherwise seem unconcerned 2—sobering of attention, slight concern expressed 3—moderate concern, relatively clear but brief change to sad/sympathetic affect 4—expressed clear facial concern and/or sympathetic vocal expression	.87
Prosocial Behavior	Attempts to help the examiner	1—no helping behavior or only after a direct prompt 2—helped after multiple indirect prompts 3—helped after one indirect prompt 4—spontaneously began helping	.95
Personal Distress	Physical/bodily signs of tension/anxiety	1—no apparent distress, child appears relaxed 2—brief, relatively minor tension or anxiety expressed, child is able to overcome quickly 3—moderate body tension/freezing or apparent anxiety, which may delay helping 4—significant distress as evident by prolonged body tension/freezing, significantly interrupts behavioral response	.95

ICC, intraclass correlations (absolute agreement, single measure).

full CSS range—from a lack of symptomatology to severe symptoms. ADOS protocols were scored by research-reliable clinicians.

Analytic Plan

Preliminary analyses examined associations between empathic responding and possible covariates. Analysis of variance models were then conducted for each empathic responding variable to assess ASD outcome group differences. Significant models were followed up by post hoc comparisons of the three outcome groups (Bonferroni-corrected $P = .017$). Finally, correlations and regression models predicting SA-CSS and RRB-CSS from empathic responding variables were conducted within the high-risk group to examine ASD-related differences in empathy on a continuous metric.

Results

For the sample as a whole, correlations among empathic responding variables (see Table III) indicated that children who responded with higher levels of visible distress to an examiner's distress displayed more empathic concern and less prosocial behavior. There was a trend-level correlation suggesting that children with higher levels of parent-reported empathic responding tended to show more observed empathic concern. These correlations were similar in magnitude when children in the ASD group were removed from the analysis (see Table III), although personal distress and empathic concern were no longer correlated in the reduced sample, indicating that inclusion of the ASD group was central to this association.

Associations between child age and IQ, and empathic responding variables were also assessed. There were no significant correlations between child age and empathic

Table III. Correlations among Study Variables

Variable	EC	PB	PD	PR
Entire sample				
Empathic concern	—	.17	.25*	.24 [†]
Prosocial behavior		—	-.42**	.14
Personal distress			—	-.04
Parent report	.19	.03		—
Age	-.09	-.16	.05	.15
Full Scale IQ			.23 [†]	.26*
No ASD only				
Empathic concern	—	.21	.13	.26 [†]
Prosocial behavior		—	-.46**	.16
Personal distress			—	-.14
Parent report				—
Age	.16	.04	.01	.13
Full Scale IQ	-.15	-.17	.19	.11

Notes. [†] $P < .10$; * $P < .05$; ** $P < .01$. $N = 70-74$ for observed variables and $N = 62-66$ for parent report (entire sample). $N = 63-64$ for observed variables and $N = 55$ for parent report in (no ASD only).

responding variables regardless of inclusion of the ASD group (see Table III); age was not included in further analyses. Children with higher IQs had higher reported empathic responding and tended to show more personal distress; IQ was included as a covariate in subsequent analyses. One-way ANOVAs were conducted to assess gender differences in empathic responding (see Table IV), none of which reached significance (empathic concern: $F(1, 71) = 3.18, P = .08$; prosocial behavior: $F(1, 72) = .00, P = .98$; personal distress: $F(1, 72) = .85, P = .36$; parent report: $F(1, 64) = 1.74, P = .19$). Given the lack of gender differences in measures of empathic responding and ASD outcome groups, gender was not included in subsequent analyses.¹

¹Including gender in an ANCOVA analysis of empathic concern yielded results equivalent to the reported analysis without gender.

Table IV. Descriptive Information by Gender and ASD Outcome

Empathic responding variable <i>M</i> (<i>SD</i>)	Male	Female
Empathic concern	2.19 (.98)	2.60 (.97)
Prosocial behavior	2.65 (1.07)	2.65 (1.28)
Personal distress	2.09 (1.07)	2.32 (1.05)
Parent report	4.83 (.89)	5.12 (.81)

Empathic responding variable <i>EMM</i> (<i>SE</i>)	Low risk/no ASD	High risk/no ASD	ASD
Empathic concern	2.42 (.19)	2.51 (.17)	1.69 (.32)
Prosocial behavior	2.44 (.22)	2.78 (.20)	2.70 (.37)
Personal distress	2.50 (.19) ^a	2.28 (.17) ^a	1.39 (.32) ^b
Parent report	4.66 (.16) ^b	5.33 (.14) ^a	4.76 (.28) ^{ab}

Notes. EMM, estimated marginal means after adjusting for IQ. Parent report is the Empathy and Prosocial Response scale from Conscience Questionnaire [Kochanska et al., 1994]. ASD, autism spectrum disorder. Groups with different superscripts significantly differ at $P = .017$ level (Bonferroni-corrected).

ANCOVA models assessed ASD outcome group differences on the four outcome measures while covarying IQ (see Fig. 1 and Table IV). Differences in empathic concern by ASD outcome group did not reach significance, $F(2, 66) = 2.72, P = .07, \text{partial } \eta^2 = .08$. Likewise, there was no difference in prosocial behavior between ASD outcome groups, $F(2, 67) = .67, P = .52, \text{partial } \eta^2 = .02$.

There was a difference in personal distress² based on ASD outcome, $F(2, 67) = 4.24, P < .05, \text{partial } \eta^2 = .11$. The ASD group showed less personal distress than the low risk/no ASD group, $F(1, 67) = 8.35, P = .005$, and the high risk/no ASD group, $F(1, 67) = 6.00, P = .017$, while the high risk/no ASD and low risk groups did not differ, $F(1, 67) = .72, P = .40$. Parent-reported empathic responding differed by ASD outcome, $F(2, 59) = 5.48, P < .01, \text{partial } \eta^2 = .16$. The high risk/no ASD group had higher levels of empathic responding than the low risk/no ASD group, $F(1, 59) = 9.73, P = .003$. The ASD group did not differ from the high risk/no ASD group, $F(1, 59) = 3.24, P = .08$, or the low risk/no ASD group, $F(1, 59) = .09, P = .77$.

Pearson's correlations quantifying zero-order associations between empathic responding and ADOS SA-CSS and RRB-CSS within the high-risk group are presented in Table V. SA-CSS was moderately negatively correlated with observed empathic concern and personal distress, as well as parent-reported empathic responding. RRB-CSS was negatively correlated only with personal distress.

We further assessed the relationship between empathic responding and ADOS severity scores in regressions controlling for IQ. A model predicting SA-CSS from observed empathic responding variables was significant, $F(4,$




Figure 1. *Significant difference between groups in post hoc analyses (Bonferroni-corrected $P = .017$), following significant ANCOVA model. Empathic concern, personal distress, and prosocial behavior were rated on a 1–4 scale based on observed behavioral responses to an examiner's accident. The parent report was the Empathy/Prosocial Response subscale from the Conscience Questionnaire [Kochanska et al., 1994]. Parents rated children's behaviors on a 1–7 scale. Estimated marginal means and standard errors (controlling for IQ) are presented.

38) = 3.87, $P = .01, R^2 = .29$; in this model, empathic concern significantly predicted SA-CSS, $\beta = -.32, P = .04$. The model predicting RRB-CSS from observed empathy was not significant, $F(4, 38) = 1.24, P = .31, R^2 = .12$, nor was the model predicting RRB-CSS from parent-reported empathic responding, $F(2, 34) = 3.15, P = .06, R^2 = .16$. The overall model predicting SA-CSS from parent-reported empathic responding was significant, $F(2, 34) = 6.64, P < .01, R^2 = .28$. However, when controlling for IQ, $\beta = -.47, P < .01$, parent report ratings were no longer associated with SA-CSS, $\beta = -.12, P = .47$.

Discussion

The current study examined observed and parent-reported empathic responding in 4- to 6-year-old

²Levene's test ($P = .047$) for the personal distress variable, indicated differences in error variances between groups. Nonparametric tests confirmed differences in personal distress between groups (Kruskal–Wallis Test, $P = .009$).

Table V. Correlations between ADOS Severity and Empathic Responding Variables

Variable	Empathic concern	Prosocial behavior	Personal distress	Parent report
ADOS severity				
SA-CSS	-.39**	-.07	-.41**	-.41*
RRB-CSS	-.17	-.04	-.29*	-.31 [†]

Notes. [†] $P < .10$; * $P < .05$; ** $P < .01$. Correlations included only children with familial risk for autism. ADOS, Autism Diagnostic Observation Schedule; SA, Social Affect; RRB, Restricted, Repetitive Behaviors; CSS, Calibrated Severity Scores. Observed variables $n = 45$, parent report $n = 39$.

children with familial risk for ASD, with and without ASD outcomes, as well as low-risk siblings. Both high-risk and low-risk siblings without ASD displayed more personal distress in response to an examiner's distress than children with ASD. Strikingly, high-risk siblings without ASD were reported by their parents to show higher levels of empathic responding than the low-risk children, while the ASD group did not differ from either group in their level of parent-reported empathic responding. Within the high-risk sibling group, level of observed empathic concern was the strongest predictor of children's Social Affect symptoms on the ADOS, although this variable did not significantly differentiate the ASD outcome groups. There was no evidence of an association between prosocial behavior and ASD outcome or symptoms. In sum, children with ASD showed a lack of emotional responsiveness but adequate helping behavior, while high-risk siblings without significant ASD symptomatology were unimpaired.

The finding that early deficits in emotional responsiveness to distress in high-risk siblings with ASD persist into preschool age supplements previous studies of high-risk siblings during the first years of life [Hutman et al., 2010; McDonald & Messinger, 2012]. As in past studies, empathy deficits were specific to ASD outcome and not familial risk status, indicating a lack of support for the involvement of (emotional) empathy deficits in the BAP. In fact, parents in this study reported higher levels of empathic responding in high-risk siblings without ASD than in low-risk siblings. Additionally, higher levels of observed and parent-reported empathic responding were associated with fewer social-communication deficits. Our findings are consistent with a separate early school-age follow-up study of high-risk siblings indicating a lack of significant difference in parent-reported affective empathy abilities, and non-significantly higher mean scores in the high-risk siblings without ASD vs. the ASD and low risk groups [Tsang et al., 2016].

It is possible that an increased capacity for empathy has a protective effect in children who did not develop ASD despite elevated familial risk, or that these children may have learned to show more empathy toward others because of experience with their affected siblings. Alternatively, parent perception may play a role. For example,

parents of high-risk siblings without ASD may perceive their child without ASD as more empathic because of implicit comparisons with the older sibling with ASD. Similarly, parents of two children with ASD may have favorably compared their younger child with ASD to an older, more affected child, which may, in part, explain why ratings of children with ASD were similar to those of low-risk children. The lack of difference in parent-reported empathic responding between the ASD and non-ASD groups may also reflect the limited inclusion of cognitive aspects of empathy in our parent report measure, the area in which deficits were most evident in previous studies examining parent-reported empathy in children with ASD [e.g., Tsang et al., 2016].

Results from this study contribute to the literature on empathic responding in preschool-aged children with ASD. Children with ASD showed different patterns of responses to an examiner's accident than children without ASD. Specifically, children with ASD showed less personal distress than high-risk and low-risk siblings without ASD. ASD outcome group differences in empathic concern did not reach significance, although empathic concern proved the strongest predictor of ADOS Social Affect scores in high-risk siblings. Together, these results indicate that children with higher levels of ASD symptoms, as indicated by both ASD diagnosis and more severe social-communication deficits, show less emotional responsiveness to an examiner's distress. The low levels of apparent personal distress in the children with ASD and the association between ASD symptoms and empathic concern are consistent with previous literature showing lower physiological responsiveness [Corona, Dissanayake, Arbelle, Wellington, & Sigman, 1998] and reduced attention and facial affect [Sigman et al., 1992] in response to adult distress in children with ASD. It was somewhat unexpected that personal distress, which is considered a less developmentally mature response to distress, best differentiated the groups. The observational measure of empathy occurred in a fairly unfamiliar lab space with a relatively unfamiliar adult. It is possible that the unfamiliarity of the situation may have differentially elicited initial anxiety or wariness in some of the non-ASD children, which they were then able to regulate to subsequently express concern or provide help.

Although ASD-related deficits in the emotional aspects of empathic responding were evident, children

with ASD were as likely to provide help to an unfamiliar adult as children without ASD. This is consistent with earlier work in high-risk siblings [McDonald & Messinger, 2012] and previous studies in children with ASD [Deschamps et al., 2014; Liebal, Colombi, Rogers, Warneken, & Tomasello, 2008], suggesting that empathic responding deficits in ASD may not be as pervasive as was once believed. These findings are in contrast with evolutionary arguments that empathy is the primary mechanism for motivating prosocial and altruistic behavior [de Waal, 2008], suggesting the possibility of a different mechanism for motivating prosocial behavior in children with ASD. Motivation for the prosocial behavior of the children with ASD may have instead come from a belief in a sense of fairness or learned responding.

Intercorrelations among empathic responding variables indicated that parent-reported empathic responding was marginally associated with empathic concern, indicating that the questionnaire may have most closely reflected individual differences in behaviors signifying concern for others. Observed empathic concern was also positively associated with personal distress, although this association was no longer significant when children with ASD were removed from the sample. This pattern of findings suggests that the children with ASD may have had an outsized effect on the correlation between personal distress and empathic concern. For children with ASD, personal distress may have been a signal that they were emotionally affected by the examiner's distress, which then may have motivated a display of empathic concern.

With regard to prosocial behavior, a negative correlation between prosocial behavior and personal distress was observed, which was unsurprising as high levels of personal distress are thought to interfere with helping behavior [Eisenberg et al., 2010]. The lack of correlation between empathic responding and prosocial behavior was not expected [e.g., Eisenberg et al., 2010]. Research suggests, however, that not all types of prosocial behavior are associated with empathy [e.g., Carlo & Randall, 2002]. While it is possible that helping behavior measured by the current task was not motivated by empathic concern, the lack of significant correlation may have been due to limited power.

Study limitations impact the interpretation of these findings and suggest avenues for future research. Although overall sample size was moderate, there were relatively few children with ASD outcomes ($n = 12$); thus, the null findings reported with respect to the ASD sample in this study must be considered with caution and may not generalize to the broader population of children with ASD. In particular, for observed empathic concern, the small ASD group size may have limited our power to detect a significant group difference, given

the trend toward lower ratings for the children with ASD ($P = .07$). Further, while we included a parent report and observational measure of empathic responding, it would be informative if future research examined children's responses to a range of social partners, including peers, and empathy-inducing situations. Along these lines, the addition of a specific measure of cognitive empathy would have allowed for a clearer comparison with previous studies [e.g., Deschamps et al., 2014; Tsang et al., 2016]. This study also did not include physiological measures that would provide information on children's internal responses to distress.

The current study provides a detailed examination of empathic responding, including parent report and an analysis of different components of empathy-related behavioral responses. This analysis indicated areas of deficit (i.e., emotional responsiveness) and areas of relative strength (i.e., helping behavior) associated with ASD outcome and symptoms. The unique sample, children with familial risk for ASD, also allowed us to investigate whether high-risk siblings without ASD showed evidence of (lower-level) empathy deficits related to the BAP. We did not find such deficits. Rather, high-risk siblings without ASD showed evidence of comparable (or even higher) levels of empathy as low-risk children. This finding, together with evidence that empathy deficits precede diagnosis and predict the severity of later autism symptoms in high-risk siblings [Hutman et al., 2010; McDonald & Messinger, 2012] suggests that empathy may be protective in high-risk siblings. Our findings highlight the importance of examining children's empathic responding as a multi-dimensional construct when examining individual differences and support empathic responding as a therapeutic target in children at risk for ASD.

Acknowledgments

The research reported in this article was supported by grants from the National Institute of Child Health and Development (R01HD057284 and R01HD047417), the National Institute of General Medical Sciences (R01GM105004), and the Marino Autism Research Institute. We thank the families who participated in this research. The authors do not have any conflicts of interest associated with this study.

References

- American Psychiatric Association. (2000). DSM-IV-TR: Diagnostic and statistical manual of mental disorders, text revision. Washington, DC: American Psychiatric Association.
- Bacon, A.L., Fein, D., Morris, R., Waterhouse, L., & Allen, D. (1998). The responses of autistic children to the distress of others. *Journal of Autism and Developmental Disorders*, 28, 129–142.

- Batson, C.D., Fultz, J., & Schoenrade, P.A. (1987). Distress and empathy: Two qualitatively distinct vicarious emotions with different motivational consequences. *Journal of Personality*, 55, 19–39.
- Carlo, G., & Randall, B.A. (2002). The development of a measure of prosocial behaviors for late adolescents. *Journal of Youth and Adolescence*, 31, 31–44.
- Coplan, R.J., & Arbeau, K.A. (2009). Peer interactions and play in early childhood. In K.H. Rubin, B. Bukkowsky, & B. Laursen (Eds.), *Handbook of peer interactions, relationships, and groups* (pp 143–161). New York, NY: Guilford Press.
- Corona, R., Dissanayake, C., Arbelle, S., Wellington, P., & Sigman, M. (1998). Is affect aversive to young children with autism? Behavioral and cardiac responses to experimenter distress. *Child Development*, 69, 1494–1502.
- Dawson, G., Toth, K., Abbott, R., Osterling, J., Munson, J., Estes, A., & Liaw, J. (2004). Early social attention impairments in autism: Social orienting, joint attention, and attention to distress. *Developmental Psychology*, 40, 271–283.
- Denham, S.A., Blair, K.A., DeMulder, E., Levitas, J., Sawyer, K., Auerbach-Major, S., & Queenan, P. (2003). Preschool emotional competence: Pathway to social competence? *Child Development*, 74, 238–256.
- Deschamps, P.K., Been, M., & Matthys, W. (2014). Empathy and empathy induced prosocial behavior in 6-and 7-year-olds with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 44, 1749–1758.
- De Waal, F.B. (2008). Putting the altruism back into altruism: The evolution of empathy. *Annual Review of Psychology*, 59, 279–300.
- Eisenberg, N., & Eggum, N.D. (2009). Empathic responding: Sympathy and personal distress. In J. Decety & W. Ickes (Eds.), *The social neuroscience of empathy*. Cambridge, MA: The MIT Press, pp 71–83.
- Eisenberg, N., Eggum, N.D., & Giunta, L.D. (2010). Empathy-related responding: Associations with prosocial behavior, aggression, and intergroup relations. *Social Issues Policy Review*, 4, 143–180.
- Eisenberg, N., Fabes, R.A., Murphy, B., Karbon, M., Smith, M., & Maszk, P. (1996). The relations of children's dispositional empathy-related responding to their emotionality, regulation, and social functioning. *Developmental Psychology*, 32, 195–209.
- Eisenberg, N., Fabes, R.A., & Spinrad, A. (2006). Prosocial development. In W. Damon & N. Eisenberg (Eds.), *Handbook of child psychology* (pp 646–718). New York, NY: Wiley.
- Gotham, K., Pickles, A., & Lord, C. (2009). Standardizing ADOS scores for a measure of severity in autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 39, 693–705.
- Grove, R., Baillie, A., Allison, C., Baron-Cohen, S., & Hoekstra, R.A. (2014). The latent structure of cognitive and emotional empathy in individuals with autism, first-degree relatives and typical individuals. *Molecular Autism*, 5, 42.
- Hoffman, M.L. (2008). Empathy and prosocial behavior. In: Lewis M., Haviland-Jones J.M., Barrett L.F., editors. *Handbook of emotions*, 3e. New York: Guilford Press, pp 440–455.
- Hus, V., Gotham, K., & Lord, C. (2014). Standardizing ADOS domain scores: Separating severity of social affect and restricted and repetitive behaviors. *Journal of Autism and Developmental Disorders*, 44, 2400–2412.
- Hutman, T., Rozga, T., DeLaurentis, A.D., Barnwell, J.M., Sugar, C.A., & Sigman, M. (2010). Response to distress in infants at risk for autism: A prospective longitudinal study. *Journal of Child Psychology and Psychiatry*, 51, 1010–1020.
- Kochanska, G., DeVet, K., Goldman, M., Murray, K., & Putnam, S.P. (1994). Maternal reports of conscience development and temperament in young children. *Child Development*, 65, 852–868.
- Liebal, K., Colombi, C., Rogers, S.J., Warneken, F., & Tomasello, M. (2008). Helping and cooperation in children with autism. *Journal of Autism and Developmental Disorders*, 38, 224–238.
- Lord, C., Risi, S., Lambrecht, L., Cook, E.H. Jr., Leventhal, B.L., DiLavore, P.C., ... Rutter, M. (2000). The Autism Diagnostic Observation Schedule—Generic: A standard measure of social and communication deficits associated with the spectrum of autism. *Journal of Autism and Developmental Disorders*, 30, 205–223.
- Lord, C., Rutter, M., & Le Couteur, A. (1994). Autism Diagnostic Interview-Revised: A revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. *Journal of Autism and Developmental Disorders*, 24, 659–685.
- McDonald, N.M., & Messinger, D.S. (2012). Empathic responding in toddlers at risk for an autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 42, 1566–1573.
- Piven, J. (2001). The broad autism phenotype: A complementary strategy for molecular genetic studies of autism. *American Journal of Human Genetics*, 105, 34–35.
- Sagi, A., & Hoffman, M.L. (1976). Empathic distress in the newborn. *Developmental Psychology*, 12, 175–176.
- Sigman, M.D., Kasari, C., Kwon, J.H., & Yirmiya, N. (1992). Responses to the negative emotions of others by autistic, mentally retarded, and normal children. *Child Development*, 63, 796–807.
- Scheeren, A.M., Koot, H.M., Mundy, P.C., Mous, L., & Begeer, S. (2013). Empathic responsiveness of children and adolescents with high-functioning autism spectrum disorder. *Autism Research*, 6, 362–371.
- Svetlova, M., Nichols, S.R., & Brownell, C.A. (2010). Toddlers' prosocial behavior: From instrumental to empathic to altruistic helping. *Child Development*, 81, 1814–1827.
- Tsang, T., Gillespie-Lynch, K., & Hutman, T. (2016). Theory of mind indexes the broader autism phenotype in siblings of children with autism at school age. *Autism Research and Treatment*, 2016, 1–13.
- Wechsler, D. (2002). *The Wechsler Preschool and Primary Scale of Intelligence, Third Edition (WPPSI-III)*. San Antonio, TX: The Psychological Corporation.
- Wellman, H.M., Cross, D., & Watson, J. (2001). Meta-analysis of theory-of-mind development: The truth about false belief. *Child Development*, 72, 655–684.
- Young, S.K., Fox, N.A., & Zahn-Waxler, C. (1999). The relations between temperament and empathy in 2-year-olds. *Developmental Psychology*, 35, 1189–1197.
- Zahn-Waxler, C., Radke-Yarrow, M., Wagner, E., & Chapman, M. (1992). Development of concern for others. *Developmental Psychology*, 28, 126–136.
- Zhou, Q., Eisenberg, N., Losoya, S.H., Fabes, R.A., Reiser, M., Guthrie, I.K., ... Shepard, S.A. (2002). The relations of parental warmth and positive expressiveness to children's empathy-related responding and social functioning: A longitudinal study. *Child Development*, 73, 893–915.